

Probabilistic Analysis of Rule 2

Jennie C. Hansen

Actuarial Math and Statistics Department
Herriot-Watt University
J.Hansen@ma.hw.ac.uk

Eric Schmutz

Department of Mathematics
Drexel University
Philadelphia, Pa. 19104
Eric.Jonathan.Schmutz@drexel.edu

Li Sheng

Department of Mathematics
Drexel University
Philadelphia, Pa. 19104
lsheng@cs.drexel.edu

February 1, 2008

Abstract

Li and Wu proposed Rule 2, a localized approximation algorithm that attempts to find a small connected dominating set in a graph. Here we study the asymptotic performance of Rule 2 on random unit disk graphs formed from n random points in an $\ell_n \times \ell_n$ square region of the plane. If $\ell_n = O(\sqrt{n/\log n})$, Rule 2 produces a dominating set whose expected size is $O(n/(\log \log n)^{3/2})$.

keywords and phrases: *coverage process, dominating set, localized algorithm, performance analysis, probabilistic analysis, Rule k , unit disk graph*

1 Introduction

Suppose random points $V_1, V_2, \dots, V_n$ are selected from a connected region $\mathcal{Q}$ in $\mathbb{R}^2$. For each i , let $D_1(V_i)$ be the unit disk centered at V_i . There is a large literature on coverage processes[18] that enables one to answer questions such as whether or not the random disks are likely to cover all of $\mathcal{Q}$, i.e. whether $\mathcal{Q} \subseteq \bigcup_{i=1}^n D_1(V_i)$. A variant question asks whether there is small *subset* of the disks whose union already covers $\mathcal{Q}$: given $k < n$, are there indices $i_1 < i_2 < \dots < i_k$ such that $\mathcal{Q} \subseteq \bigcup_{j=1}^k D_1(V_{i_j})$. For this variant, there are several interesting ways to modify the meaning of “coverage.” For example: is there a small subset of the disks whose union is connected and contains all n points $V_1, V_2, \dots, V_n$ (but not necessarily all of $\mathcal{Q}$)? These questions are a bit vague, but specific examples arise naturally in connection with probabilistic models for wireless networks. In particular, they are central to the probabilistic analysis of Rule 2 in this paper.

Rule 2 is a well known algorithm that was proposed by Wu and Li [31] as a means of increasing the efficiency of routing in ad hoc wireless networks. To describe the algorithm and a probabilistic model, we need some graph theoretic terminology. A *unit disk graph* has for its vertex set $\mathcal{V}$ a finite set of points in $\mathbb{R}^2$. Given the vertex set $\mathcal{V}$, the edge set $\mathcal{E}$ is determined as follows: an *undirected* edge $e \in \mathcal{E}$ connects vertices $u, v \in \mathcal{V}$ (and in this case we say that u and v are adjacent) iff $d(u, v)$, the Euclidean distance between them, is less than one. Unit disk graphs have been used by many authors as mathematical models for the interconnections between nodes in a wireless network, and random unit disk graphs have been used as probabilistic models for these networks [8],[12], [15],[16],[17],[23],[24]. A *dominating set* in any graph $G = (\mathcal{V}, \mathcal{E})$ is a subset $\mathcal{C} \subseteq \mathcal{V}$ such that every vertex $v \in \mathcal{V}$ either is in the set $\mathcal{C}$, or is adjacent to a vertex in $\mathcal{C}$. We say $\mathcal{C}$ is a *connected dominating set* if $\mathcal{C}$ is a dominating set and the subgraph induced by $\mathcal{C}$ is connected. Of course it is not possible for G to have a connected dominating set if G itself is not connected. We use the acronym “CDS” for a dominating set $\mathcal{C}$ such that the subgraph induced by $\mathcal{C}$ has the same number of components that G has. This paper deals with a *random* unit disk graph model, $\mathcal{G}_n$, which is connected with asymptotic probability one. Thus any CDS for $\mathcal{G}_n$ will also be connected with high probability. We assume that each vertex has a unique identifier taken from a totally ordered set. For convenience, when $|\mathcal{V}| = n$, we will use the numbers $1, 2, \dots, n$ as IDs, and will number the vertices accordingly. If v_i is any vertex (with ID i), define the neighborhood $\mathcal{N}(v_i)$ to be the set consisting of v_i and any vertices in $\mathcal{V}$ that are adjacent to v_i . The CDS constructed by the Rule 2 algorithm is denoted $\mathcal{C}(\mathcal{V})$, and its cardinality is $C(\mathcal{V}) = |\mathcal{C}(\mathcal{V})|$. The elements of $\mathcal{C}(\mathcal{V})$ are called “gateway nodes”. $\mathcal{C}(\mathcal{V})$ consists of all vertices $v_i \in \mathcal{V}$ that are not excluded under the following version of Rule 2:

Rule 2: Vertex v_i is excluded from $\mathcal{C}(\mathcal{V})$ iff $\mathcal{N}(v_i)$ contains at least one set of two vertices v_{i_1}, v_{i_2} such that

- $i_1 > i_2 > i$ and
- $\mathcal{N}(v_i) \subseteq \mathcal{N}(v_{i_1}) \cup \mathcal{N}(v_{i_2})$ and
- v_{i_1} is adjacent to v_{i_2} .

Wu and Li showed that this algorithm produces a CDS. They also conjectured, based on simulation data, that it is effective in the sense that it selects a CDS that is small relative to n “in the average case”. In this paper we treat the analysis of Rule 2 mathematically by considering its performance when it is applied to a random unit disk graph $\mathcal{G}_n$. Specifically, let $\ell_1 \leq \ell_2 \leq \dots$ be a sequence of real numbers such that $\ell_n = O(\sqrt{n/\log n})$ as $n \rightarrow \infty$, but $\ell_n \geq \log n$ for all n . Let $\mathcal{Q}_n$ be an $\ell_n \times \ell_n$ square region in $\mathbb{R}^2$. Select n points $V_1, V_2, \dots, V_n$ independently and uniform randomly from an $\mathcal{Q}_n$, and use these n points as the vertex set for a unit disk graph $\mathcal{G}_n$. With this probabilistic model, the size of the Rule 2 dominating set is a random variable. We prove asymptotic estimates for the expected size of the Rule 2 dominating set. The proof involves some interesting problems in elementary geometry and geometric probability.

2 A Geometric Lemma

As observed in [20], a unit disk centered at a point o cannot be completely covered with two unit disks having centers at points u and w ($u \neq o \neq w$): $(D_1(u) \cup D_1(w))^c \cap D_1(o) \neq \emptyset$. One might infer that a typical vertex o is not likely to be pruned under Rule 2 because no two points in $\mathcal{N}(o)$ will cover all the vertices in $\mathcal{N}(o)$. This reasoning suggests that Rule 2 will be ineffective. But such reasoning is not sound. Typically there are points u and w that cover all but a negligible fraction of the disk centered at o . The uncovered region is small enough so that it usually does not include any nodes. A more precise version of this statement is proved in the next section, but first we need to look carefully at the area of regions such as $(D_1(u) \cup D_1(w))^c \cap D_1(o)$. In particular, we need Lemma 1, which is the main result in this section.

To state Lemma 1 we adopt some notation. Throughout this section $b > 1$ will be a parameter and in terms of b we let $L = \lfloor b^{1/3}(\log b)^2 \rfloor$, $\delta = \frac{1}{\sqrt[3]{b \log b}}$, and $\theta_b = \pi/L$. We fix $o = (x_o, y_o) \in \mathbb{R}^2$ and for any $r > 0$, let $D_r(o)$ be the closed disk centered at o with radius r . We are going to partition the small disk $D_\delta(o)$ into $2L$ sectors as follows. Choose a new coordinate system centered at o , and for $0 \leq i < L$, let Q_i be the sector consisting of those points $(x, y) = (r \cos \theta, r \sin \theta)$ whose polar coordinates satisfy $0 < r \leq \delta$ and $(i - \frac{1}{2})\theta_b \leq \theta \leq (i + \frac{1}{2})\theta_b$. Similarly let R_i be the sector that is obtained by reflecting Q_i about o , namely the points with $0 < r < \delta$ and $(i - \frac{1}{2})\theta_b < \theta - \pi < (i + \frac{1}{2})\theta_b$. The analysis of Rule 2 depends on a geometric lemma about these sectors. For any i , and any points $q_i \in Q_i, u_i \in R_i$, let $X(q_i, u_i)$ be the area of $(D_1(q_i) \cup D_1(u_i))^c \cap D_1(o)$, i.e. the area of the omitted region in $D_1(o)$ that is not covered by $(D_1(q_i) \cup D_1(u_i))$. Let $\tilde{q}_i$ and $\tilde{u}_i$ be the extreme points whose polar coordinates are respectively $(r, \theta) = (\delta, (i - \frac{1}{2})\theta_b)$ and $(r, \theta) = (\delta, (i + \frac{1}{2})\theta_b + \pi)$. We prove:

Lemma 1 *There is a uniform constant $C > 0$ such that, for $0 \leq i < L$, and for all $q_i \in Q_i, u_i \in R_i$, we have $X(q_i, u_i) \leq X(\tilde{q}_i, \tilde{u}_i) \leq \frac{C}{b \log^3 b}$.*

Proof.

We prove four facts which together imply Lemma 1. In the first fact, we observe that omitted area $X(q, u)$ gets larger if we move one (or both) of the two points q, u away from the origin along a radial line.

Fact 1 *Let q_1, q_2 and u_1, u_2 be four points in $D_1(o)$ such that q_1 lies on the line segment $\overline{o, q_2}$ and u_1 lies on the line segment $\overline{o, u_2}$. Then $X(q_2, u_2) \geq X(q_1, u_1)$.*

Proof. It suffices to show that $D_1(q_2) \cap D_1(o) \subseteq D_1(q_1) \cap D_1(o)$ and that $D_1(u_2) \cap D_1(o) \subseteq D_1(u_1) \cap D_1(o)$. Suppose $p \in D_1(q_2) \cap D_1(o)$. Since q_1 lies on the line segment from o to q_2 , we have $d(q_1, p) \leq \max(d(o, p), d(q_2, p)) \leq 1$. Hence $p \in D_1(q_1) \cap D_1(o)$. By a similar same argument, $D_1(q_2) \cap D_1(o) \subseteq D_1(q_1) \cap D_1(o)$. □

Fact 2 *Let a, b be the two points where the circles $\partial D_1(p), \partial D_1(q)$ intersect. Then, $\overline{a, b} \perp \overline{p, q}$, and the two line segments $\overline{a, b}$ and $\overline{p, q}$ intersect at their midpoints.*

Proof. This follows immediately from the fact that $d(p, a) = d(p, b) = d(q, a) = d(q, b) = 1$. □

Fact 3 *Let o_1, o_2 be two points on the circle $x^2 + y^2 = \delta^2$. Then, $X(o_1, o_2)$ is a decreasing function of $\angle o_1 o o_2$.*

Proof. For convenience, we will use polar coordinates. Without loss of generality, let o_1 be the point with polar coordinates $(r_{o_1}, \phi_{o_1}) = (\delta, \pi)$. Let o_2 be an arbitrary point on the circle with the polar coordinates (δ, ϕ_2) . By symmetry, we only need to consider the case when o_2 is in the first or second quadrant; we may, without loss of generality, assume that $0 \leq \phi_2 \leq \pi$. We will show that $X(o_1, o_2)$ is an increasing function of ϕ_2 , then the result follows from the fact that $\angle o_1 o o_2 = \pi - \phi_2$.

Let a_1, b_1 be the two points where the circles $\partial D_1(o_1)$ and $\partial D_1(o)$ intersect, with a_1 in the second quadrant and b_1 in the third quadrant.

Let o^* be a point on the circle $x^2 + y^2 = \delta^2$ so that $\partial D_1(o^*)$ meets with both $\partial D_1(o)$ and $\partial D_1(o_1)$ at a_1 . Let b^*, d^* be the other intersection points of $\partial D_1(o^*)$ with $\partial D_1(o)$ and $\partial D_1(o_1)$, respectively. For convenience, let's denote ϕ_{o^*} by ϕ^* . Figure 1 illustrates the position of $\partial D_1(o_1), \partial D(o)$, and $\partial D_1(o^*)$ and their intersections.




Figure 1: The position of the circle $\partial D_1(o^*)$

As in the proof of Fact 2, we have $\overline{a_1, d^*} \perp \overline{o_1, o^*}$, $\overline{a_1, b^*} \perp \overline{o, o^*}$. Notice also that o is on the line segment $\overline{a_1, d^*}$. So,

$$\angle b^* a_1 o = \angle o o^* o_1 = \angle o^* o_1 o = \frac{\phi^*}{2}. \quad (1)$$

It follows that

$$0 < \phi^*/2 < \pi/2, \text{ and, } \sin \frac{\phi^*}{2} = \frac{\delta}{2} \quad (2)$$

Now, for the point o_2 with polar coordinates (δ, ϕ_2) , let a_2, b_2 denote the two points where $\partial D_1(o_2)$ and $\partial D_1(o)$ intersect, and let c_2, d_2 denote the two points where $\partial D_1(o_2)$ and $\partial D_1(o_1)$ intersect. There are two cases to consider: $\phi_2 \leq \phi^*$, and $\phi_2 \geq \phi^*$.

Case 1. $\phi_2 \leq \phi^*$.




Figure 2: The case when $\phi_2 \leq \phi^*$

Notice that a_1, b_1 partitions the circle $\partial D_1(o)$ into two arcs: the right section and the left section. When, $\phi_2 \leq \phi^*$, as illustrated in Figure 2, a_2, b_2 are both on the right section of the circle $\partial D_1(o)$ between a_1, b_1 . Similarly, c_2, d_2 are both on the right section of the circle $\partial D_1(o_1)$ between a_1, b_1 . Clearly,

$$X(o_1, o_2) = B_1 - (B_2 - B_3) = B_1 - B_2 + B_3,$$

where

- $B_1 = \text{area}(D_1(o_1)^c \cap D_1(o))$
- $B_2 = \text{area}(D_1(o) \cap D_1(o_2))$
- $B_3 = \text{area}(D_1(o_1) \cap D_1(o_2))$, the shaded area in Figure 2

Notice that B_3 is the only area that depends on ϕ_2 . We shall now give an expression for B_3 .

Let's denote $\angle c_2 o_1 o_2 = y$. Since $\angle o_2 o_1 o = \frac{\phi_2}{2}$, we have

$$0 < y < \frac{\pi}{2}, \text{ and, } \cos y = \delta \cos \frac{\phi_2}{2} \quad (3)$$

By symmetry, one can see that the shaded region is partitioned equally by the line $\overline{c_2, d_2}$. So,

$$B_3 = 2\left(\frac{2y}{2\pi}\pi - \frac{1}{2}(2 \sin y)(\cos y)\right) = 2y - \sin 2y.$$

Here, the first term is the area of the sector $D_1(o_1)$ that extends from c_2 to d_2 , and the second term is the area of the triangle (c_2, o_1, d_2) .

From the above two equations, we have

$$\frac{dX(o_1, o_2)}{d\phi_2} = \frac{dB_3}{d\phi_2} = \frac{dB_3}{dy} \cdot \frac{dy}{d\phi_2} = (1 - \cos 2y) \cdot \frac{\delta \sin \frac{\phi_2}{2}}{2 \sin y} > 0.$$

Here the last inequality follows from the fact that $0 < \frac{\phi_2}{2}, y < \frac{\pi}{2}$. Thus $X(o_1, o_2)$ is an increasing function in ϕ_2 .

Case 2. $\phi_2 > \phi^*$.




Figure 3: The case when $\phi_2 > \phi^*$

One can see from Figure 3 that

$$X(o_1, o_2) = B_1 - (B_2 - B_3) = B_1 - B_2 + B_3$$

Where B_1, B_2 are defined the same as those in the case 1, but

$B_3 = \text{area}(D_1(o_1) \cap D_1(o_2) \cap D_1(o))$, the shaded area in Figure 3

Again, B_3 is the only area that depends on ϕ_2 . We will now give an expression of B_3 .

We show first that $\angle c_2 o a_1 = \angle a_2 o c_2$ by showing that $\phi_{c_2} - \phi_{a_1} = \phi_{a_2} - \phi_{c_2}$. Then, it follows that B_3 is split in half by the line segment c_2, d_2 .

From Figure 1, one can see that

$$\phi_{a_1} = \phi^* + \left(\frac{\pi}{2} - \angle b^* a_1 o\right) = \phi^* + \left(\frac{\pi}{2} - \frac{\phi^*}{2}\right) = \frac{\pi}{2} + \frac{\phi^*}{2} \quad (4)$$

To find ϕ_{a_2} , observe that, as in the proof of Fact 2, we have $\overline{a_2, b_2} \perp \overline{o, o_2}$. So, $\sin \angle b_2 a_2 o = \frac{\delta}{2}$. Comparing with (2), we see that $\sin \angle b_2 a_2 o = \sin \frac{\phi^*}{2}$. This implies that $\angle b_2 a_2 o = \frac{\phi^*}{2}$. Thus,

$$\phi_{a_2} = \phi_2 + \left(\frac{\pi}{2} - \angle b_2 a_2 o\right) = \phi_2 + \left(\frac{\pi}{2} - \frac{\phi^*}{2}\right) \quad (5)$$

Now, for c_2 , using the fact that $\overline{c_2, o} \perp \overline{o_1, o_2}$,

$$\phi_{c_2} = \pi - \left(\frac{\pi}{2} - \angle o_2 o_1 o\right) = \pi - \left(\frac{\pi}{2} - \frac{\phi_2}{2}\right) = \frac{\pi}{2} + \frac{\phi_2}{2} \quad (6)$$

It follows that $\phi_{c_2} - \phi_{a_1} = \phi_{a_2} - \phi_{c_2} = \frac{\phi_2}{2} - \frac{\phi^*}{2}$. Now, using that the circle $\partial D_1(o_1)$ in the polar system is

$$r = \sqrt{1 - \delta^2 \sin^2 \phi} - \delta \cos \phi$$

and that

$$\phi_{d_2} = -(\pi - \phi_{c_2}) = -(\frac{\pi}{2} - \frac{\phi_2}{2}) \quad (7)$$

we get

$$\begin{aligned} B_3 &= 2 \left(\int_{-(\frac{\pi}{2} - \frac{\phi_2}{2})}^{\frac{\pi}{2} + \frac{\phi_2^*}{2}} \int_0^{\sqrt{1 - \delta^2 \sin^2 \phi - \delta \cos \phi}} r dr d\phi + \frac{\frac{\phi_2}{2} - \frac{\phi_2^*}{2}}{2\pi} \cdot \pi \right) \\ &= \int_{-(\frac{\pi}{2} - \frac{\phi_2}{2})}^{\frac{\pi}{2} + \frac{\phi_2^*}{2}} 1 - \delta^2 \sin^2 \phi + \delta^2 \cos^2 \phi - 2\delta \cos \phi \sqrt{1 - \delta^2 \sin^2 \phi} d\phi + \frac{\phi_2 - \phi_2^*}{2} \end{aligned}$$

Thus,

$$\begin{aligned} \frac{dX(o_1, o_2)}{d\phi_2} = \frac{dB_3}{d\phi_2} &= -\frac{1}{2} [1 - \delta^2 \sin^2(-\frac{\pi}{2} + \frac{\phi_2}{2}) + \delta^2 \cos^2(-\frac{\pi}{2} + \frac{\phi_2}{2}) \\ &\quad - 2\delta \cos(-\frac{\pi}{2} + \frac{\phi_2}{2}) \sqrt{1 - \delta^2 \sin^2(-\frac{\pi}{2} + \frac{\phi_2}{2})}] + \frac{1}{2} \\ &= \frac{1}{2} [\delta^2 \cos^2 \frac{\phi_2}{2} - \delta^2 \sin^2 \frac{\phi_2}{2} + 2\delta \sin \frac{\phi_2}{2} \sqrt{1 - \delta^2 \cos^2 \frac{\phi_2}{2}}] \\ &= \frac{1}{2} [-(\delta \sin \frac{\phi_2}{2} - \sqrt{1 - \delta^2 \cos^2 \frac{\phi_2}{2}})^2 + 1] \\ &\geq 0 \end{aligned}$$

The last inequality follows because $0 \leq \delta \sin \frac{\phi_2}{2} \leq 1$, $0 \leq \sqrt{1 - \delta^2 \cos^2 \frac{\phi_2}{2}} \leq 1$, and thus $(\delta \sin \frac{\phi_2}{2} - \sqrt{1 - \delta^2 \cos^2 \frac{\phi_2}{2}})^2 < 1$. $\square$

Fact 4 *Uniformly for all i , we have $X(\tilde{q}_i, \tilde{u}_i) = O(\frac{1}{b \log^3 b})$.*

Proof. Without loss of generality, let $i = 0$ and $v = (0, 0)$. To simplify notation, define $x_b = \delta \cos(-\frac{1}{2}\theta_b)$, $y_b = \delta \sin(-\frac{1}{2}\theta_b)$. Let (ξ, η) be the point in the first quadrant where the circles $x^2 + y^2 = 1$ and $(x - x_b)^2 + (y - y_b)^2 = 1$ meet. Then

$$\begin{aligned} X(\tilde{q}_0, \tilde{u}_0) &\leq 4 \int_0^\xi \sqrt{1 - x^2} - (y_b + \sqrt{1 - (x - x_b)^2}) dx \\ &= -4y_b \xi + 4 \int_0^\xi \frac{-2xx_b + x_b^2}{\sqrt{1 - x^2} + \sqrt{1 - (x - x_b)^2}} dx \end{aligned}$$

Hence we have

$$X(\tilde{q}_0, \tilde{u}_0) = O(\xi y_b) + O(x_b \xi^2) + O(x_b^2 \xi). \quad (8)$$

Note that $x_b^2 + y_b^2 = \delta^2 = \frac{1}{b^{2/3} \log^2 b}$, that $\xi^2 + \eta^2 = 1$, that $(\xi - x_b)^2 + (\eta - y_b)^2 = 1$, that $x_b = \delta(1 + O(\theta_b^2))$, and that $y_b = \frac{-\delta\theta_b}{2}(1 + O(\theta_b^2))$. Combining these equations, we get $\xi = O(\delta)$. Putting this estimate back into (8), we get

$$X(\tilde{q}_0, \tilde{u}_0) = O(\frac{1}{b \log^3 b}). \quad (9)$$

□

In the analysis of Rule 2 it is necessary to consider vertices in $\mathcal{G}_n$ which are close to the boundary of the square $\mathcal{Q}_n$. For this reason we define, for $o \in \mathbb{R}_+^2$, the “truncated unit disk” $\hat{D}_1(o) := D_1(o) \cap \mathbb{R}_+^2$ and we note that $\hat{D}_1(o) \subseteq D_1(o)$, and $\hat{D}_1(o) = D_1(o)$ iff $x_o, y_o \geq 1$. Then for L and δ as defined above, we have the following corollary to Lemma 1:

Corollary 2 *There is a uniform constant $C > 0$ such that, for all $o \in \mathbb{R}_+^2$ such that $D_\delta(o) \subseteq \mathbb{R}_+^2$, for $0 \leq i < L$, and for all $q_i \in Q_i, u_i \in R_i$, we have $\hat{X}(q_i, u_i) \leq X(\tilde{q}_i, \tilde{u}_i) \leq \frac{C}{b \log^3 b}$, where $\hat{X}(q, u)$ is the area of $(D_1(q) \cap D_1(u))^c \cap \hat{D}_1(o)$.*

Proof. Clearly $\hat{X}(q_i, u_i) \leq X(q_i, u_i)$ since $\hat{D}_1(o) \subseteq D_1(o)$. So the result follows from Lemma 1 (since $\tilde{q}_i, \tilde{u}_i \in D_\delta(o) \subseteq \mathbb{R}_+^2$). □

3 Local Coverage by Two Discs

Recall that under Rule 2 a vertex v_i is excluded from $\mathcal{C}(\mathcal{V})$ if there are two adjacent vertices, $v_{i_1}, v_{i_2} \in \mathcal{N}(v_i)$, with higher IDs than v_i which also ‘cover’ v_i , i.e. $\mathcal{N}(v_i) \subseteq \mathcal{N}(v_{i_1}) \cup \mathcal{N}(v_{i_2})$. In the analysis of Rule 2 we will distinguish vertices in $\mathcal{N}(v_i)$ with higher ID than v_i by coloring them blue; all other vertices in $\mathcal{N}(v_i)$ are colored white. With this in mind, we consider in this section a two-colored random unit disk graph and prove a local coverage result.

Let w and b be positive integers such that $w < b(\log b)^2$ and, as before, let $L = \lfloor b^{1/3}(\log b)^{3/2} \rfloor$ and $\delta = \frac{1}{b^{1/3} \log b}$. Fix $o \in \mathbb{R}_+^2$ such that $D_\delta(o) \subseteq \mathbb{R}_+^2$ and select $w+b$ points independently and uniform randomly from the truncated disk $\hat{D}_1(o)$. Color the first w points white, and the remaining b points blue. Form a random (improperly colored) unit disk graph $\hat{\mathcal{H}}_{w,b}$ by putting an edge between two of the $w+b$ colored points iff the distance between them is one or less. Our goal in this section is to prove that, with high probability, $\hat{\mathcal{H}}_{w,b}$ contains a dominating set consisting of two blue vertices that are adjacent to each other.

For $0 \leq i < L$, let Q_i, R_i denote the sectors of $D_\delta(o)$ as defined in the previous section and let $N(Q_i), N(R_i)$ respectively be the number of blue vertices of $\hat{\mathcal{H}}_{w,b}$ that lie in Q_i and R_i . Let $\tau_b = \sum_{i=0}^{L-1} I_i$ where, in this section only, the $I_i = 1$ if and only if $N(R_i) = N(Q_i) = 1$ (and otherwise $I_i = 0$.) We note that the distribution of τ_b depends on the position of o and we indicate this dependence by using the notation $\Pr_o(\tau_b \in \cdot)$. Provided o is not too close to the boundary of $\mathbb{R}_+^2$, we can obtain uniform bounds on the tail of the distribution of τ_b :

Lemma 3 $\Pr_o\left(\tau_b < \frac{b^{1/3}}{16 \log^6 b}\right) = O\left(\frac{\log^6 b}{b^{1/3}}\right)$ uniformly for all $o \in \mathbb{R}_+^2$ such that $D_\delta(o) \subseteq \mathbb{R}_+^2$.

Proof. Let $|\hat{D}_1(o)|$ denote the area of $\hat{D}_1(o)$, let $\hat{\lambda} = \hat{\lambda}(o) = \frac{\pi}{|\hat{D}_1(o)|}$, and define

$$\hat{p} = \frac{\text{Area}(Q_i)}{|\hat{D}_1(o)|} = \frac{\pi \delta^2 / 2L}{|\hat{D}_1(o)|} = \frac{\hat{\lambda}}{2b \log^4 b} \left(1 + O\left(\frac{1}{b^{1/3} \log^2 b}\right)\right). \quad (10)$$

The expected value of I_i depends on o :

$$E_o(I_i) = b(b-1)\hat{p}^2(1-2\hat{p})^{b-2} = \frac{\hat{\lambda}^2}{4\log^8 b} \left(1 + O\left(\frac{1}{\log^4 b}\right)\right). \quad (11)$$

Hence

$$E_o(\tau_b) = LE_o(I_i) = \frac{b^{1/3}\hat{\lambda}^2}{4\log^6 b} \left(1 + O\left(\frac{1}{\log^4 b}\right)\right). \quad (12)$$

We likewise have, for $i \neq j$,

$$E_o(I_i I_j) = b(b-1)(b-2)(b-3)\hat{p}^4(1-4\hat{p})^{b-4} = \frac{\hat{\lambda}^4(o)}{16\log^{16} b} \left(1 + O\left(\frac{1}{\log^4 b}\right)\right). \quad (13)$$

Note that

$$\pi \geq |\hat{D}_1(o)| \geq \frac{\pi}{4}, \quad (14)$$

and therefore

$$1 \leq \hat{\lambda}(o) \leq 4. \quad (15)$$

Therefore we have uniformly for all $o \in \mathfrak{R}_+^2$ such that $D_\delta(o) \subseteq \mathfrak{R}_+^2$

$$Var(\tau_b) = O\left(\frac{b^{1/3}}{\log^6 b}\right). \quad (16)$$

Observe that

$$\Pr_o\left(\tau_b < \frac{b^{1/3}}{16\log^6 b}\right) \leq \Pr_o\left(\tau_b \leq \frac{1}{2}E_o(\tau_b)\right) \leq \Pr_o\left(|\tau_b - E_o(\tau_b)| > \frac{1}{2}E_o(\tau_b)\right). \quad (17)$$

The lemma now follows from (16), (17) and Chebyshev's inequality. $\square$

Recall our assumptions that $w < b(\log b)^{3/2}$, that $\delta = \frac{1}{b^{1/3}\log b}$, and that $x_o, y_o \geq \delta$. With these assumptions, we have:

Theorem 4 *There is a constant $c > 0$, independent of the position of o , such that with probability at least $1 - \frac{c}{(\log b)^{3/2}}$, the random graph $\hat{\mathcal{H}}_{w,b}$ has a connected dominating set that consists of two blue vertices in $D_\delta(o)$.*

Proof.

Let $\mathcal{T}_b \subseteq \{0, 1, 2, 3, \dots, L-1\}$ be the random subset of indices such that $i \in \mathcal{T}_b$ iff $N(Q_i) = N(R_i) = 1$. If $\mathcal{T}_b \neq \emptyset$, define $Y = \min \mathcal{T}_b$ to be the smallest of the indices in $\mathcal{T}_b$; otherwise, if $\mathcal{T}_b = \emptyset$, set $Y = -1$.

Define the random variable X_b as follows: If $\tau_b = |\mathcal{T}_b| = 0$ then $X_b = 0$; otherwise, if $\mathcal{T}_b = \{i_1, i_2, \dots, i_{\tau_b}\}$ and $i_1 < i_2 < \dots < i_{\tau_b}$, then $X_b = 1$ iff $Q_{i_1} \cup R_{i_1}$ contains a blue connected dominating set for $\hat{\mathcal{H}}_{w,b}$.

Let $\mathcal{B} = \{g_1, g_2, \dots, g_b\}$ be the set of blue nodes, selected independently and uniform randomly from $\hat{D}_1(o)$. Define $\mathcal{Z} = \mathcal{B} \cap D_\delta(o)$ to be set of blue points that fall near the origin o , and let $Z = |\mathcal{Z}|$ be the number of these points. Then

$$\Pr_o(X_b = 0) \leq \Pr_o\left(X_b = 0, \tau_b \neq 0, Z \leq \frac{2\hat{\lambda}b^{1/3}}{(\log b)^2}\right) + \Pr_o(\tau_b = 0) + \Pr_o\left(Z > \frac{2\hat{\lambda}b^{1/3}}{(\log b)^2}\right). \quad (18)$$

Note that Z has a binomial distribution: $Z \stackrel{d}{=} \text{Bin}(b, \hat{\lambda}\delta^2)$ where $\hat{\lambda}$ is as defined in the proof of Lemma 3. If $\beta = \frac{2\hat{\lambda}b^{1/3}}{(\log b)^2}$, then by Chernoff's inequality,

$$\Pr_o(Z \geq \beta) \leq \exp(-b^{1/3}/4(\log b)^2). \quad (19)$$

By Lemma 3, $\Pr_o(\tau_b = 0) = O(\frac{\log^6 b}{b^{1/3}})$. Therefore

$$\Pr_o(X_b = 0) \leq \Pr_o(X_b = 0, \tau_b \neq 0, Z \leq \beta) + O(\frac{\log^6 b}{b^{1/3}}). \quad (20)$$

Now we decompose the first term on the right side of (20) according to the value of Y .

$$\Pr_o(X_b = 0, \tau_b \neq 0, Z \leq \beta) = \sum_{k=0}^{L-1} \Pr_o(X_b = 0 | Y = k, Z \leq \beta) \Pr_o(Y = k, Z \leq \beta). \quad (21)$$

(The redundant condition $\tau_b \neq 0$ need not be included on the right side of (21) because it a consequence of the condition $Y \geq 0$.) We have

$$\Pr_o(X_b = 0 | Y = k, Z \leq \beta) = \sum_S \Pr_o(X_b = 0 | \mathcal{Z} = S, Y = k) \Pr_o(\mathcal{Z} = S | Y = k, Z \leq \beta) \quad (22)$$

where the sum is over subsets $S \subseteq [b]$ such that $2 \leq |S| \leq \beta$.

$$\Pr(X_b = 0 | \mathcal{Z} = S, Y = k) = 1 - \Pr(X_b = 1 | \mathcal{Z} = S, Y = k), \quad (23)$$

so it is enough to find a lower bound for $\Pr(X_b = 1 | \mathcal{Z} = S, Y = k)$.

To simplify notation, let $\gamma = X(\tilde{q}_0, \tilde{u}_0)$, and recall that $\gamma = O(\frac{1}{b \log^3 b})$. In this section of the paper, define $|D_\delta(o)| = \frac{\pi}{b^{2/3}(\log b)^2}$ to be the area of the disk $D_\delta(o)$, and let $|\hat{D}_1(o)| = \text{Area}(\hat{D}_1(o))$. An important observation is that, once we have specified $b - |S|$ = the number of blue points that fall *outside* $D_\delta(o)$, the locations in $D_\delta(o)^c \cap \hat{D}_1(o)$ of these $b - |S|$ points are independent of the locations of the $|S|$ blue points *in* $D_\delta(o)$, and are also independent of the locations of the white points. Hence

$$\Pr_o(X_b = 1 | \mathcal{Z} = S, Y = k) \geq \frac{(1 - \frac{|D_\delta(o)|}{|\hat{D}_1(o)|} - \frac{\gamma}{|D_1(o)|})^{b-|S|}}{(1 - \frac{|D_\delta(o)|}{|\hat{D}_1(o)|})^{b-|S|}} \left(1 - \frac{\gamma}{|\hat{D}_1(o)|}\right)^w \quad (24)$$

$$\geq \left(1 - \frac{C}{b(\log b)^3}\right)^{b-|S|+w} \quad (25)$$

for some constant C that is independent of o . With our assumption $w < b(\log b)^{3/2}$ we get, for all sufficiently large b , the lower bound

$$\Pr_o(X_b = 1 | \mathcal{Z} = S, Y = k) \geq \left(1 - \frac{C'}{b(\log b)^3}\right)^{b(\log b)^{3/2}} \geq 1 - \frac{C''}{(\log b)^{3/2}} \quad (26)$$

for some constants C' and C'' which are independent of $\mathcal{Z}, Y$, and o . Hence

$$\Pr_o(X_b = 0) \leq \frac{c}{(\log b)^{3/2}} \quad (27)$$

for some constant c that is *independent of the point o* . □

4 Analysis of Rule 2

Let U be the number of nodes that become non-gateways when Rule 2 is applied to the random graph $\mathcal{G}_n$: $U = \sum_i I_i$ where (in this section) the indicator variable $I_i = 1$ iff the node with ID i becomes a non-gateway under Rule 2. Assume that there is a positive constant $\bar{c}$ such that, for all $n > 1$, $\log n \leq \ell_n \leq \bar{c}\sqrt{\frac{n}{\log n}}$. Let $\xi_n = \frac{\alpha_n}{\ell_n^2}$, where $\langle \alpha_n \rangle$ is any sequence of real numbers satisfying the following three conditions:

- $\alpha_n = o(n)$ as $n \rightarrow \infty$.
- $\xi_n = \frac{\alpha_n}{\ell_n^2} \rightarrow \infty$ as $n \rightarrow \infty$.
- For all sufficiently large n , $\frac{16n}{\log^{3/2} \xi_n} < \alpha_n$.

For example, if $\ell_n = \Theta(\sqrt{n/\log n})$, then the sequence $\alpha_n = \frac{32n}{(\log \log n)^{3/2}}$ satisfies the three conditions. On the other hand, if $\ell_n = \Theta((n/\log n)^t)$ for some fixed positive $t < 1/2$, then $\alpha_n = \frac{n}{\log n}$ satisfies the three conditions above. With these three assumptions, our goal is to prove

Theorem 5 $E(U) \geq n - O(\alpha_n)$.

Proof. The idea of the proof is to use Theorem 4 to bound the probability that a typical vertex V_i is pruned by Rule 2. In this case the blue vertices correspond to nodes in $D_1(V_i)$ with IDs *higher* than i , and the white vertices correspond to nodes in $D_1(V_i)$ with lower IDs. Let $r = \frac{1}{\log^{3/2} \xi_n}$, and let $\mathcal{A}_i$ be the event that $D_r(V_i) \subseteq \mathcal{Q}_n$. Then

$$\Pr(\mathcal{A}_i) = \frac{(\ell_n - 2r)^2}{\ell_n^2} \geq 1 - \frac{4r}{\ell_n}. \quad (28)$$

Let $\hat{D}_1(V_i) = D_1(V_i) \cap \mathcal{Q}_n$ be the set of points in $\mathcal{Q}_n$ whose distance from V_i is one or less, and let $|\hat{D}_1(V_i)|$ be the area of $\hat{D}_1(V_i)$. Let $\rho_i^{(b)}$ denote the number

of nodes in $\hat{D}_1(V_i)$ having a label that is larger than i , and let $\rho_i^{(w)}$ be the number of nodes in $\hat{D}_1(V_i)$ having a label that is smaller than i . Then, given the location of the i 'th vertex V_i , $\rho_i^{(b)}$ has a Binomial($n-i, \frac{|\hat{D}_1(V_i)|}{\ell_n^2}$) distribution. Define $\mu_b = \mu_b(i)$ to be the expected value of $\rho_i^{(b)}$ given the location of the i 'th point:

$$\mu_b = E(\rho_i^{(b)}|V_i) = \frac{(n-i)|\hat{D}_1(V_i)|}{\ell_n^2}. \quad (29)$$

Similarly $\rho_i^{(w)}$ has a Binomial($i-1, \frac{|\hat{D}_1(V_i)|}{\ell_n^2}$) distribution, and we define $\mu_w = \mu_w(i)$ to be the expected value:

$$\mu_w = E(\rho_i^{(w)}|V_i) = \frac{(i-1)|\hat{D}_1(V_i)|}{\ell_n^2}. \quad (30)$$

If $\mathcal{A}_i$ occurs, then by Chebyshev's inequality,

$$\Pr(|\rho_i^{(b)} - \mu_b(i)| < \frac{\mu_b}{2} | \mathcal{A}_i) \geq 1 - \frac{16\ell_n^2}{n-i}, \quad (31)$$

and similarly for $\rho_i^{(w)}$.

If we let $\mathcal{D}_i$ be the event that *both* of the inequalities $|\rho_i^{(b)} - \mu_b(i)| < \frac{\mu_b}{2}$ and $|\rho_i^{(w)} - \mu_w(i)| < \frac{\mu_w}{2}$ are satisfied, then

$$\Pr(\mathcal{D}_i | \mathcal{A}_i) \geq 1 - \frac{16\ell_n^2}{n-i} - \frac{16\ell_n^2}{i-1}. \quad (32)$$

Combining (32) and (28), we get

$$\Pr(\mathcal{D}_i \cap \mathcal{A}_i) \geq \left(1 - \frac{16\ell_n^2}{n-i} - \frac{16\ell_n^2}{i-1}\right) \left(1 - \frac{4r}{\ell_n}\right). \quad (33)$$

Now let $\lambda_n = n - \alpha_n$, then clearly

$$E(U) \geq \sum_{i=\alpha_n}^{\lambda_n} \Pr(I_i = 1) \geq \sum_{i=\alpha_n}^{\lambda_n} \Pr(I_i = 1 | \mathcal{D}_i \cap \mathcal{A}_i) \Pr(\mathcal{D}_i \cap \mathcal{A}_i) \quad (34)$$

To obtain a lower bound for the right hand side of inequality (34), we prove

Lemma 6 *There is a constant $\tilde{c} > 0$ such that for all sufficiently large n and all $\alpha_n \leq i < \lambda_n$, $\Pr(I_i = 1 | \mathcal{D}_i \cap \mathcal{A}_i) \geq 1 - \frac{\tilde{c}}{(\log \xi_n)^{3/2}}$.*

Proof. We begin by noting that *given* the event $\mathcal{D}_i \cap \mathcal{A}_i$ and $\alpha_n \leq i < \lambda_n = n - \alpha_n$, we have

$$\rho_i^{(w)} < \frac{3}{2}\mu_w(i) = \frac{3(i-1)|\hat{D}_1(V_i)|}{2\ell_n^2} \leq \frac{3\pi n}{2\ell_n^2}. \quad (35)$$

Similarly

$$\rho_i^{(b)} > \frac{1}{2}\mu_b(i) = \frac{(n-i)|\hat{D}_1(V_i)|}{2\ell_n^2} > \frac{\alpha_n\pi}{8\ell_n^2} = \frac{\xi_n\pi}{8} \quad (36)$$

It follows from inequalities (35) and (36) and from the conditions on the sequences $\langle \xi_n \rangle$ and $\langle \alpha_n \rangle$ that, *given* $\mathcal{D}_i \cap \mathcal{A}_i$ and $\alpha_n \leq i < \lambda_n$,

$$\rho_i^{(b)}(\log \rho_i^{(b)})^{3/2} \geq \rho_i^{(w)}. \quad (37)$$

Next we consider the conditional probability $\Pr(I_i = 1 | \rho_i^{(b)}, \rho_i^{(w)}, V_i, \mathcal{D}_i \cap \mathcal{A}_i)$ where the values of $\rho_i^{(b)}$ and $\rho_i^{(w)}$ and the location of V_i are consistent with the event $\mathcal{D}_i \cap \mathcal{A}_i$. In this case, it follows from inequality (36) that

$$\delta(\rho_i^{(b)}) := \frac{1}{(\rho_i^{(b)})^{1/3} \log(\rho_i^{(b)})} \leq \frac{1}{(\xi_n/3)^{1/3} \log(\xi_n/3)} \leq \frac{1}{(\log(\xi_n))^{3/2}} = r. \quad (38)$$

Since the event $\mathcal{A}_i$ implies $D_r(V_i) \subseteq \mathfrak{R}_+^2$, it follows from (38) that $D_{\delta(\rho_i^{(b)})}(V_i) \subseteq \mathfrak{R}_+^2$. Finally, it follows from Theorem 4 that for some fixed positive constant $\tilde{c}$

$$\Pr(I_i = 1 | \rho_i^{(b)}, \rho_i^{(w)}, V_i, \mathcal{D}_i \cap \mathcal{A}_i) \geq 1 - \frac{c}{(\log(\rho_i^{(b)}))^{3/2}} \geq 1 - \frac{\tilde{c}}{(\log(\xi_i^{(b)}))^{3/2}} \quad (39)$$

for all sufficiently large n and all $\alpha_n \leq i < \lambda_n$. The lemma now follows from (39). $\square$

Recall that $\lambda_n = n - \alpha_n$, that $\alpha_n = o(n)$, that $\xi_n = \frac{\alpha_n}{\ell_n^2} \rightarrow \infty$ as $n \rightarrow \infty$, and that for all sufficiently large n , $\alpha_n > \frac{16n}{(\log \xi_n)^{3/2}}$. So it follows from Lemma 6 and (33) and (34), that

$$E(U) \geq n - 2\alpha_n + o(\alpha_n).$$

$\square$

5 Discussion

In this final section, assume $\ell_n = \Theta((\frac{n}{\log n})^t)$ for some fixed positive $t \leq \frac{1}{2}$. For all sufficiently large n , the expected size of the Rule 2 dominating set is at least $\ell_n^2/4$ (See Theorem 5 of [17]). There is a gap between this lower bound and the $O(\alpha_n)$ upper bound in Theorem 5. For example, when $t = 1/2$, the lower and upper bounds for the expected size of the Rule 2 dominating set are respectively $\Theta(n/\log n)$ and $\Theta(n/(\log \log n)^{3/2})$. For $t < 1/2$ the gap is even wider: the lower and upper bounds are respectively $\Theta((\frac{n}{\log n})^{2t})$ and $\Theta(\frac{n}{\log n})$. We conjecture that, in fact, the expected size of the Rule 2 dominating set is $\Theta(\ell_n^2)$.

Acknowledgement We thank Jie Wu for introducing us to this problem. Professor Wu made helpful suggestions regarding an early version of this paper.

References

- [1] C.Adjih,P.Jacquet, L.Viennot,Computing Connected Dominating Sets with Multipoint Relays, *INRIA RR-4597* (2002).
- [2] K.M.Alzoubi,P.J.Wan,and O.Frieder, Distributed Heuristics for Connected Dominating Sets in Wireless Ad Hoc Networks, *Journal of Communications and Networks*,**4** (1) (2002) 1–8.
- [3] M.J.B.Appel and R.P.Russo, The minimum vertex degree of a graph on uniform points in $[0, 1]^d$, *Advances in Appl.Probab* **29** (3) (1997) 582–594.
- [4] E.Baccelli and P.Jacquet, Flooding Techniques in Mobile Ad Hoc Networks, *INRIA RR-5002* (2003).
- [5] T.Camp and B.Williams, Comparison of Broadcasting Techniques for Mobile Ad Hoc Networks, Proceedings of the 3’rd ACM International Symposium on Mobile Ad Hoc Networking and Computing, MobiHoc 2002 (2002) 194–205.
- [6] M.Cardei,X.Cheng,X.Cheng, D.Du,Connected Domination in Multihop Ad Hoc Wireless Networks, *JCIS* (2002) 251–255.
- [7] B. Chen, K. Jamieson, H. Balakrishnan, R. Morris , Span: An Energy-Efficient Coordination Algorithm for Topology Maintenance in Ad Hoc Wireless Networks *Proc. of the 6th ACM MOBICOM Conf.*, Rome, Italy, July 2001.
- [8] B.N. Clark,C.J. Colburn, and D.J.Johnson, Unit Disk Graphs,*Discrete Mathematics* **86**(1-3) (1990) 165–177.
- [9] B.Das and V. Bharghavan Routing in Ad-Hoc Networks Using Minimum Connected Dominating Sets, International Conference on Communications **1** (1997) 376–380.
- [10] F.Dai and J.Wu, An Extended Localized Algorithm for Connected Dominating Set Formation in Ad Hoc Wireless Networks, *IEEE Transactions on Parallel and Distributed Systems*, **15** (10) 2004.
- [11] F.Dai and J.Wu, Performance analysis of broadcast protocols in ad hoc networks based on self pruning, *IEEE Trans.on Parallel and Distributed Systems* **15** (11) (2004).
- [12] E.N.Gilbert, Random Plane Networks, *J.Soc.Indust.Appl.Math.* **9** (1961) 533.

- [13] A.Godbole and B.Wielund, On the Domination Number of a Random Graph, *Electronic Journal of Combinatorics* **8** #R37 (2001)
- [14] S.Guha and S.Khuller, Approximation algorithms for connected dominating sets, *Algorithmica* **20** (4) (1998) 374–387.
- [15] P.Gupta and P.R.Kumar, “Critical power for asymptotic connectivity in wireless networks”, in *Stochastic Analysis, Control, Optimization and Applications*, Birkhauser (1999) 547–566.
- [16] W.K.Hale, Frequency Assignment: Theory and Applications, *Proc. IEEE* **68** (1980) 1497–1514.
- [17] J.C.Hansen and E.Schmutz, The Expected size of the Rule k dominating set, *submitted*.
- [18] P.Hall, *Introduction to the Theory of Coverage Processes*, Wiley (1988).
- [19] P.Jacquet, Analytical Results on Connected Dominating Sets in Mobile Ad Hoc Networks, *INRIA RR-5173* (2004).
- [20] P.Jacquet, A.Laouiti, P.Minet, L.Viennot, Performance of Multipoint Relaying in Ad Hoc Mobile Routing Protocols, In “Networking 2002” *Lecture Notes in Computer Science* **2345** (2002) 387–398.
- [21] D.Lichtenstein, Planar formulae and their uses, *SIAM J.Comput.* **11**(2) (1982) 329–343.
- [22] M.V.Marathe, H.Breu, H.B.Hunt, S.S.Ravi, and D.J.Rosenkrantz, Simple Heuristics for Unit Disk Graphs, *Networks* **25** no.2 (1995) 59–68.
- [23] C.McDiarmid, Discrete mathematics and radio channel assignment, *Recent Advances in algorithms and combinatorics*, 27–63, CMS Books Math (2003).
- [24] C.McDiarmid, Random channel assignment in the plane, *Random Structures and Algorithms* **22** (2) 187–212.
- [25] W.Peng and X.Lu, On the reduction of broadcast redundancy in mobile ad hoc networks, *Proceedings of the 1st ACM International Symposium on Mobile Ad Hoc Networking* (2000) 129–130.
- [26] *Random Geometric Graphs*, Oxford Studies in Probability **5**, Oxford University Press, (2003) ISBN 0-19-850626-0.
- [27] R.Sivakumar, B.Das, and V. Bharghavan, , Spine-based routing in ad hoc networks, *Cluster Computing* **1** (2) (1998) 237–248.
- [28] Solomon, *Geometric Probability*, *CBMS-NSF Conference Series* **28** SIAM, 1978.

- [29] I. Stojmenovic, M. Seddigh, J. Zunic, Dominating sets and neighbor elimination based broadcasting algorithms in wireless networks, *IEEE Transactions on Parallel and Distributed Systems*, Vol. **13**, No. **1**, (2002), 14-25.
- [30] Y.C.Tseng, S.Y.Ni, Y.S.Chen, J.P.Sheu, The broadcast storm problem in a mobile ad hoc network, *Wireless Networks* **8(2-3)** (2002) 153-167.
- [31] J. Wu and H.Li, On calculating connected dominating set for efficient routing in ad hoc wireless networks, *Workshop on Discrete Algorithms and Methods for MOBILE Computing and Communications* (1999) 7-14.
- [32] A.C.C.Yao, On constructing spanning trees in k dimensional spaces and related problems, *SIAM J.Computing* **11 (4)** (1982) 721-736.